arXiv:1211.3637v2 [math.GT] 20 Dec 2012

TAUT FOLIATIONS IN SURFACE BUNDLES WITH MULTIPLE PUNCTURES

TEJAS KALELKAR AND RACHEL ROBERTS

ABSTRACT. Let M be a fibered 3-manifold with multiple boundary components. We show that the fiber structure of M can be perturbed to yield transversely oriented taut foliations realizing all rational multislopes in some open neighborhood of the multislope of the fiber.

1. INTRODUCTION

A (codimension-1) foliation $\mathcal{F}$ of a 3-manifold M is a disjoint union of injectively immersed 2-manifolds (called leaves) such that $(M, \mathcal{F})$ looks locally like $(\mathbb{R}^3, \mathbb{R}^2 \times I)$. For a precise definition see for example [14]. Every 3-manifold has a foliation [14, 29]; however the foliations obtained from this general construction have torus leaves that bound a solid torus foliated by planes. Such a solid torus is called a Reeb component [23]. As every 3-manifold has such a foliation, it is necessary to look for foliations without Reeb components in order to attribute topological properties to the manifold. Taut foliations provide an important class of foliations without Reeb components. A taut foliation is a codimension-1 foliation of a 3-manifold for which there exists a closed curve that transversely intersects each leaf of the foliation. Taut foliations have been extensively studied by Thurston [27] and Gabai [4, 5, 6].

The existence of a taut foliation in a closed orientable 3-manifold has important topological consequences for the manifold. For example, if M is a closed, orientable 3-manifold that has a taut foliation with no sphere leaves then M is covered by $\mathbb{R}^3$ [19], M is irreducible [25] and has an infinite fundamental group [9]. In fact, its fundamental group acts nontrivially on interesting 1-dimensional objects (see, for example, [27, 1] and [19, 22]). Moreover, taut foliations can be perturbed to interesting contact structures [3] and hence can be used to obtain Heegaard-Floer information [18].

Thurston [26] gave a way to represent fibers of a fibration of M over S^1 as lattice points in the Thurston norm unit ball. This focused attention on the study of fibrations of M that are near a given fibration. Roberts [20, 21] perturbed fibrations in a compact 3-manifold with one boundary component to taut foliations that are 'nearby' in terms of their boundary behaviours. This construction was used by Honda, Kazez and Matić [10] to prove that a large class of contact structures are not only tight, but symplectically semi-fillable.

We extend Roberts' theorem [20] for compact fibered 3-manifolds with one boundary component to the multiple boundary component case.

Definition 1.1. Let F be a compact oriented surface of positive genus and with nonempty boundary consisting of k components. Let h be an orientation preserving homeomorphism of F which fixes each boundary component pointwise. Let $M = F \times I/(x,1) \sim (h(x),0)$ and denote the k (toral) boundary components of ∂M by T^1, T^2, ..., T^k.

We use the given surface bundle structure on M to fix a coordinate system on each of the boundary tori as follows. (See, for example, Section 9.G of [24] for a definition and description of coordinate system.) For each j we choose as longitude $\lambda^j = \partial F \cap T^j$, with orientation inherited from the orientation of F. For each j, we then fix as meridian μ_j an oriented simple closed curve dual to λ_j. Although, as described in [21], it is possible to use the homeomorphism h to uniquely specify such simple closed curves μ_j, we choose not to do so in this paper as all theorem statements are independent of the choice of meridional multislope.

We say a taut foliation $\mathcal{F}$ in M *realizes boundary multislope* $(m^j)_{j=1}^k$ if for each j, $1 \leq j \leq k$, $\mathcal{F} \cap T^j$ is a foliation of T^j of slope m^j in the chosen coordinate system of T^j.

The main result we shall prove in this paper is the following:

Theorem 1.2. *Given an orientable, fibered compact 3-manifold, a fibration with fiber surface of positive genus can be perturbed to yield transversely oriented taut foliations realizing a neighborhood of rational boundary multislopes about the boundary multislope of the fibration.*

As an immediate corollary for closed manifolds we therefore have:

Corollary 1.3. *Let* $\widehat{M}(r^j)$ *be the closed manifold obtained from* M *by Dehn filling* M *along the multicurve with rational multislope* $(r^j)_{j=1}^k$. *When* (r^j) *is sufficiently close to the multislope of the fibration,* $\widehat{M}(r^j)$ *also has a transversely oriented taut foliation.*

In a surface bundle with multiple boundaries, a priori there is no reason to suspect that perturbing the slope of the foliation on one boundary component would not alter boundary slopes on the other components. In other words, there is no reason to expect the various intervals of realizable boundary slopes about the slope of a fiber to be independent of each other. Our result obtains such intervals around the boundary multislope of the fiber.



FIGURE 1. Local model of a standard spine



FIGURE 2. Local model of a branched surface



FIGURE 3. Oriented spine to oriented branched surface

2. PRELIMINARIES

In this section we introduce basic definitions and fix conventions used in the rest of the paper.

2.1. **Spines and Branched surfaces.**

Definition 2.1. A *standard spine* [2] is a space X locally modeled on one of the spaces of Figure 1. The *critical locus* of X is the 1-complex of points of X where the spine is not locally a manifold.

Definition 2.2. A *branched surface* ([28]; see also [16, 17]) is a space B locally modeled on the spaces of Figure 2. The *branching locus* L of B is the 1-complex of points of B where B is not locally a manifold. The components of $B \setminus L$ are called the *sectors* of B. The points where L is not locally a manifold are called *double points* of L. They correspond to points of intersection of three sectors of B.

A standard spine X together with an orientation in a neighborhood of the critical locus determines a branched surface B in the sense illustrated in Figure 3.

Example 2.3. Let $F_0 := F \times \{0\}$ be a fiber of $M = F \times I/(x,1) \sim (h(x),0)$. Let α_i, $1 \leq i \leq k$, be pairwise disjoint, properly embedded arcs



FIGURE 4. A sink disk

in F_0, and set $D_i = \alpha_i \times I$ in M. Isotope the image arcs $h(\alpha_i)$ as necessary so that the intersection $(\cup\alpha_i) \cap (\cup_i h(\alpha_i))$ is transverse and minimal. Assign an orientation to F and to each D_i. Then $X = F_0 \cup_i D_i$ is a transversely oriented spine. We will denote by $B =< F; \cup_i D_i >$ the transversely oriented branched surface associated with $X = F_0 \cup_i D_i$.

Similarly, $\langle \cup_i F_i; \cup_{i,j} D_i^j \rangle$ will denote the transversely oriented branched surface associated to the transversely oriented standard spine

$$X = F_0 \cup F_1 \cup ... \cup F_{n-1} \cup_{i,j} D_i^j$$

where $F_i = F \times \{i/n\}$ and $D_i^j = \alpha_i^j \times [\frac{i}{n}, \frac{i+1}{n}]$ for some set of arcs α_i^j properly embedded in F so that the intersection $(\cup_j \alpha_{i-1}^j) \cap (\cup_j \alpha_i^j)$ is transverse and minimal.

Definition 2.4. A lamination carried by a branched surface B in M is a closed subset λ of an I-fibered regular neighborhood $N(B)$ of B, such that λ is a disjoint union of injectively immersed 2-manifolds (called leaves) that intersect the I-fibers of $N(B)$ transversely.

2.2. **Laminar branched surfaces.**

In [12, 13], Li introduces the fundamental notions of sink disk and half sink disk.

Definition 2.5. [12, 13] Let B be a branched surface in a 3-manifold M. Let L be the branching locus of B and let X denote the union of double points of L. Associate to each component of $L \setminus X$ a vector (in B) pointing in the direction of the cusp. A *sink disk* is a disk branch sector D of B for which the branch direction of each component of $(L \setminus X) \cap \overline{D}$ points into D. A *half sink disk* is a sink disk which has nonempty intersection with ∂M.

Sink disks and half sink disks play a key role in Li's notion of laminar branched surface.

Definition 2.6. (Definition 1.3, [12]) Let D_1 and D_2 be the two disk components of the horizontal boundary of a $D^2 \times I$ region in $M \setminus int(N(B))$. If the projection $\pi : N(B) \to B$ restricted to the interior of $D_1 \cup D_2$ is injective, i.e, the intersection of any I-fiber of $N(B)$ with $int(D_1) \cup int(D_2)$ is either empty or a single point, then we say that $\pi(D_1 \cup D_2)$ forms a *trivial bubble* in B.

Definition 2.7. (Definition 1.4, [12]) A branched surface B in a closed 3-manifold M is called a *laminar* branched surface if it satisfies the following conditions:

1. $\partial_h N(B)$ is incompressible in $M \setminus int(N(B))$, no component of $\partial_h N(B)$ is a sphere and $M \setminus B$ is irreducible.
2. There is no monogon in $M \setminus int(N(B))$; i.e., no disk $D \subset M \setminus int(N(B))$ with $\partial D = D \cap N(B) = \alpha \cup \beta$, where $\alpha \subset \partial_v N(B)$ is in an interval fiber of $\partial_v N(B)$ and $\beta \subset \partial_h N(B)$
3. There is no Reeb component; i.e., B does not carry a torus that bounds a solid torus in M.
4. B has no trivial bubbles.
5. B has no sink disk or half sink disk.

Gabai and Oertel introduced essential branched surfaces in [7] and proved that any lamination fully carried by an essential branched surface is an essential lamination and conversely any essential lamination is fully carried by an essential branched surface. In practice, to check if a manifold has an essential lamination, the tricky part often is to verify that a candidate branched surface does in fact fully carry a lamination. Li [12] uses laminar branched surfaces to relax this requirement and prove the following:

Theorem 2.8. *(Theorem 1,* [12]*) Suppose M is a closed and orientable 3-manifold. Then*

(a) *Every laminar branched surface in M fully carries an essential lamination.*
(b) *Any essential lamination in M that is not a lamination by planes is fully carried by a laminar branched surface.*

In [13], Li notices that if a branched surface has no half sink disk, then it can be arbitrarily split in a neighborhood of its boundary train track without introducing any sink disk (or half sink disk). He is therefore able to conclude the following.

Theorem 2.9. *(Theorem 2.2,* [13]*) Let M be an irreducible and orientable 3-manifold whose boundary is a union of incompressible tori. Suppose B is a laminar branched surface and $\partial M \setminus \partial B$ is a union of bigons. Then, for any multislope $(s_1, ..., s_k) \in (\mathbb{Q} \cup \{\infty\})^k$ that can be realized by the train track ∂B, if B does not carry a torus that bounds a solid torus in $\widehat{M}(s_1, ..., s_k)$, then B fully carries a lamination $\lambda_{(s_1,...,s_k)}$ whose boundary consists of the multislope $(s_1, ..., s_k)$ and $\lambda_{(s_1,...,s_k)}$ can be extended to an essential lamination in $\widehat{M}(s_1, ..., s_k)$.*



FIGURE 5. A Parallel tuple (α^i) on the surface F

We note that in [13], Li states Theorem 2.2 only for the case that ∂M is connected. However, as Li has observed and is easily seen, his proof extends immediately to the case that ∂M consists of multiple toral boundary components. Key is the fact that splitting B open, to a branched surface B' say, in a neighborhood of its boundary so that $\partial B'$ consists of multislopes $(s_1, ..., s_k)$, does not introduce sink disks. Therefore, capping B' off to $\widehat{B'}$ yields a laminar branched surface in $\widehat{M}(s_1, ..., s_k)$.

2.3. **Good oriented sequence of arcs.** In this section we introduce some definitions that will be used in the rest of the paper.

Definition 2.10. Let $(\alpha^1, ..., \alpha^k)$ be a tuple of pairwise disjoint simple arcs properly embedded in F with $\partial\alpha^j \subset T^j$. Such a tuple will be called *parallel* if $F \setminus \{\alpha^1, ..., \alpha^k\}$ has k components, $k-1$ of which are annuli $\{A^j\}$ with $\partial A^j \supset \{\alpha^j, \alpha^{j+1}\}$ and one of which is a surface S of genus $g-1$ with $\partial S \supset \{\alpha^1, \alpha^k\}$. Furthermore all α^j are oriented in parallel, i.e., the orientation of ∂A^j agrees with $\{-\alpha^j, \alpha^{j+1}\}$ and the orientation of ∂S agrees with $\{-\alpha^k, \alpha^1\}$. Note that, in particular, each α^j is non-separating. See Figure 5 for an example of a parallel tuple.

Definition 2.11. A pair of tuples $(\alpha^i)_{i=1...k}$ and $(\beta^j)_{j=1...k}$ will be called *good* if both are parallel tuples and α^i and β^j have exactly one (interior) point of intersection when $i \neq j$ while α^i is disjoint from β^j when $i = j$.

A sequence of parallel tuples

$$\sigma = ((\alpha_0^1, \alpha_0^2, ..., \alpha_0^k), (\alpha_1^1, \alpha_1^2, ..., \alpha_1^k), ..., (\alpha_n^1, \alpha_n^2, ..., \alpha_n^k))$$

also shortened to

$$((\alpha_0^j), (\alpha_1^j), ..., (\alpha_n^j))$$

or

$$(\alpha_0^j) \xrightarrow{\sigma} (\alpha_n^j)$$

will be called *good* if for each fixed j, $0 \leq j \leq k$, the pair $((\alpha_i^j), (\alpha_{i+1}^j))$ is good.



(a) (b)

FIGURE 6. A pair of arcs (α, β) in position (a) is called negatively oriented, while a pair (α, β) in position (b) is called positively oriented

Definition 2.12. We say a good pair $((\alpha^j), (\beta^j))$ is *positively oriented* if for each $j \in \{1, ..., k\}$ a neighborhood of the j-th boundary component in F is as shown in Figure 6 (b). Similarly we say a good pair $((\alpha^j), (\beta^j))$ is *negatively oriented* if for each $j \in \{1, ..., k\}$ a neighborhood of the j-th boundary component in F is as shown in Fig 6 (a).

We say a good sequence $\sigma = ((\alpha_0^j), (\alpha_1^j), ..., (\alpha_n^j))$ is positively oriented if each pair $((\alpha_i^j), (\alpha_{i+1}^j))$ is positively oriented. Similarly we say $\sigma = ((\alpha_1^j), (\alpha_2^j), ..., (\alpha_n^j))$ is negatively oriented if each pair $((\alpha_i^j), (\alpha_{i+1}^j))$ is negatively oriented. We say the sequence σ is oriented if it is positively or negatively oriented. See Figure 7 for an example of a negatively oriented good pair in F.

2.4. **Preferred generators.** Let

$$\mathcal{H}_{g,k} = \{\eta_1, \eta_2, ..., \eta_{2g-2+k,}, \gamma_{12}, \gamma_{24}, \gamma_{46}, \gamma_{68}, ..., \gamma_{2g-4,2g-2}, \beta, \beta_1, \beta_2, ..., \beta_{g-1}, \delta_1, \delta_2, ..., \delta_{k-1}\}$$

be the curves on F as shown in Figure 8. Then combining Proposition 1 and Theorem 1 of Gervais [8] the mapping class group $MCG(F, \partial F)$ of F (fixing boundary) is generated by Dehn twists about curves in $\mathcal{H}_{g,k}$.

Theorem (Gervais)**.** *The mapping class group $MCG(F, \partial F)$ of F is generated by Dehn twists about the curves in $\mathcal{H}_{g,k}$.*

As Dehn twists about δ_i are isotopic to the identity via an isotopy that does not fix the boundary, we have the following obvious corollary:

Corollary 2.13. *The mapping class group $MCG(F)$ of F (not fixing the boundary pointwise) is generated by Dehn twists about the curves in*

$$\mathcal{H}'_{g,k} = \mathcal{H}_{g,k} \setminus \{\delta_1, ..., \delta_{k-1}\}$$



FIGURE 7. Neighborhood of F with a good negatively oriented pair $((\alpha^j), (\beta^j))$ in the oriented spine X

3. MAIN THEOREM

Definition 3.1. Let $(\alpha^1, ..., \alpha^k)$ be a parallel tuple in F. Orient F so that the normal vector $\hat{n}$ induced by the orientation of M points in the direction of increasing $t \in [0, 1]$. Let $D^j = \alpha^j \times [0, 1]$ in M_h with the orientation induced by orientations of α^j and F; i.e., if v^j is tangent to α^j then $(v^j, \hat{n})$ gives the orientation of D^j. Let $X = F \cup_j D^j$ be an oriented standard spine and $B_\alpha =< F; \cup_j D^j >$ the transversely oriented branched surface associated with X.

Notice that the multislope of the fibration is $\bar{0}$. In order to prove Theorem 1.2 we shall prove the following:

Theorem 3.2. *There is an open neighborhood $\mathcal{U}$ of $\bar{0} \in \mathbb{R}^k$ such that for each point $(m^1, ..., m^k) \in \mathcal{U} \cap \mathbb{Q}^k$, there exists a lamination carried by B_α with boundary multislope (m^j). These laminations extend to taut foliations which also intersect the boundary in foliations with multislope (m^j).*

This gives us the following corollary for closed manifolds.

Corollary 3.3. *Let $\widehat{M}(r^j)$ denote the closed manifold obtained from M by a Dehn filling along a multicurve with rational multislope $(r^j)_{j=1}^k$. For each tuple $(r^j) \in \mathcal{U} \cap \mathbb{Q}^k$, the closed manifold $\widehat{M}(r^j)$ also has a transversely oriented taut foliation.*



FIGURE 8. Generators of the Mapping Class Group

The proof of Theorem 3.2 is outlined below with details worked out in the mentioned lemmas.

Proof. In Lemma 3.4 we shall show that there is a good positively oriented sequence $(\alpha_0^j) \to (h^{-1}(\alpha_0^j))$, or equivalently from $(h(\alpha_n^j)) \to (\alpha_n^j)$. In Lemma 3.6 we shall show that whenever there exists such a positive sequence there is a splitting of the branched surface B_α to a branched surface B_σ that is laminar and that therefore carries laminations realizing every multislope in some open neighborhood of $\bar{0} \in \mathbb{R}^k$. And finally in Lemma 3.8 we show that these laminations extend to taut foliations on all of M. $\square$

Lemma 3.4. *Let (α^j) be a parallel tuple in F and let $h \in Aut^+(F)$. Then there is a good positively oriented sequence $(\alpha^j) \xrightarrow{\sigma} (h(\alpha^j))$.*

Proof. By Corollary 2.13 to Gervais' Theorem, $h \sim h_m h_{m-1} ... h_2 h_1$ for twists h_i about curves in $\mathcal{H}'_{g,k}$. Set $h' = h_m h_{m-1} ... h_2 h_1$, and notice that $M_h = M_{h'}$.

By changing the handle decomposition of F as necessary, we may assume that the parallel tuple (α^j) is as shown in Figure 8. Let b denote the Dehn twist about $\beta \in \mathcal{H}'_{g,k}$. Notice that any h_i in the factorization of h' is either b, b^{-1}, or a twist about a curve disjoint from all components of α^j. Thus $((\alpha^j), (h_i(\alpha^j))$ is either a good positive pair, a good negative pair, or a pair of equal tuples.

Now if $((\alpha^j), (\beta^j))$ is a good pair then so is $((h_i(\alpha^j)), (h_i(\beta^j)))$; therefore each of the pairs
$((\alpha^j), (h_m(\alpha^j)), ((h_m(\alpha^j)), (h_m h_{m-1}(\alpha^j))), ((h_m h_{m-1}(\alpha^j)), (h_m h_{m-1}$ $h_{m-2}(\alpha^j)))$, ..., $((h_m h_{m-1} ... h_2(\alpha^j)), (h_m h_{m-1} ... h_2 h_1(\alpha^j) = h(\alpha^j)))$
is either a good oriented pair or a pair of equal tuples.

If at least one of the h_i is b or b^{-1} then ignoring the equal tuples, we get a good oriented sequence $((\alpha^j_0), (\alpha^j_1), ..., (\alpha^j_{n-1}), (\alpha^j_n) = h((\alpha^j_0)))$ or $(\alpha^j) \xrightarrow{\sigma} (h(\alpha^j))$ as required. The length of this sequence is equal to the number of times h_i equals b or b^{-1}, i.e, $n = n_+ + n_-$, where n_+ is the sum of the positive powers of b in this expression of h' and n_- is the magnitude of the sum of negative powers of b.

If none of the h_i are Dehn twists about β then $(\alpha^j) = (h(\alpha^j))$. In this case, $\sigma = ((\alpha^j), (b(\alpha^j)), (b^{-1}b(\alpha^j) = (\alpha^j)))$ is a good oriented sequence.

If $((\alpha^j), (\beta^j))$ is a positively oriented good pair then $((\alpha^j)$,$(-\beta^j)$,$(-\alpha^j)$, $(\beta^j))$ is a negatively oriented good sequence. Performing n_- such substitutions we get a positively oriented good sequence $(\alpha^j) \xrightarrow{\sigma} (h(\alpha^j))$. □

Definition 3.5. Let $\sigma = (h(\alpha^j_n) = \alpha^j_0, \alpha^j_1, ..., \alpha^j_{n-1}, \alpha^j_n)$ be a good oriented sequence. Let $F_i = F \times \{\frac{i}{n}\}$ for $0 \leq i < n$ and let $D^j_i = \alpha^j_i \times [\frac{i}{n}, \frac{i+1}{n}]$, for $0 \leq i < n$, in M_h. Let $X = (\cup_i F_i) \cup (\cup_{i,j} D^j_i)$ and orient F_i and D^j_i as in Definition 3.1. Define $B_\sigma =< \cup_i F_i; \cup_{i,j} D^j_i >$ as the associated branched surface. Figure 7 shows the neighborhood of F in X while Figure 9 shows a neighborhood of F in the associated branched surface.

Lemma 3.6. *Let $\sigma = (h(\alpha^j_n) = \alpha^j_0, \alpha^j_1, ..., \alpha^j_{n-1}, \alpha^j_n)$ be a good oriented sequence in F and B_σ the associated branched surface in M_h. Then B_σ has no sink disk or half sink disk.*

Proof. As the sequence σ is good and oriented for each fixed i, the tuple of arcs (α^j_i) is parallel and $|\alpha^j_i \cap \alpha^k_{i-1}| = \delta^k_j$, so a neighborhood of F_i in B_σ is as shown in Figure 9.

The sectors of B_σ consist of disks $D^j_i = \alpha^j_i \times [\frac{i}{n}, \frac{i+1}{n}]$ and components of $F_i \setminus \{\alpha^j_i \cup \alpha^j_{i-1}\}_{j=1...k}$. As F_{i-1} and F_i both have a co-orientation in the direction of increasing t for $(x, t) \in M_h$, so for any orientation of D^j_i, ∂D^j_i is the union of two arcs in ∂M_h, together with one arc with the direction of the cusp pointing into the disk and one arc with the direction of the cusp pointing outwards. Similarly, as α^j_i and α^{j+1}_i are oriented in parallel, each disk component of $F_i \setminus \{\alpha^j_i, \alpha^j_{i-1}\}_{j=1...k}$ has a boundary arc with cusp direction pointing outwards. Therefore no branch sector in B_σ is a sink disk or a half sink disk. □

Remark 3.7. Notice that $B_\sigma =< \cup_i F_i; \cup_{i,j} D^j_i >$ is a splitting (see [17]) of the original branched surface $B_\alpha =< F; \cup_j D^j >$ and, equivalently, B_σ



FIGURE 9. Neighborhood of one of the fibres in the branched surface B.

collapses to B_α. So in particular, laminations carried by B_σ are also carried by B_α.

Now consider the train tracks $\tau^j = B_\sigma \cap T^j$. Focus on one of the τ^j. For some choice of meridian μ_0^j. Recall that we fixed a coordinate system (λ^j, μ^j) on T^j. For simplicity of exposition, we now make a second choice μ_0^j of meridian. This choice is dictated by the form of τ^j; namely, we choose μ_0^j to be disjoint from the disks D_i^j so that τ^j has the form shown in Figure 10. Notice that there is a change of coordinates homeomorphism taking slopes in terms to the coordinate system (λ^j, μ_0^j) to slopes in terms of the coordinate system (λ^j, μ^j). Since λ^j is unchanged, this homeomorphism takes an open interval about 0 to an open interval about 0. Assign to τ^j the measure determined by weights x, y shown in Figure 10. In terms of the coordinate system (λ^j, μ_0^j), τ^j carries all slopes realizable by

$$\frac{x-y}{n(1+y)}$$



FIGURE 10. The weighted boundary train track when $n = 4$

for some $x, y > 0$. Therefore, in terms of the coordinate system (λ^j, μ_0^j), τ^j carries all slopes in $(-\frac{1}{n}, \infty)$. Converting to the coordinate system (λ^j, μ^j), τ^j carries all slopes in some open neighborhood of 0. Repeat for all j. By Theorem 2.9, we see that the branched surface B_σ carries laminations $\lambda_{(\bar{x},\bar{y})}$ realizing multislopes $(\frac{x_1-y_1}{n(1+y_1)}, \frac{x_2-y_2}{n(1+y_2)}, ..., \frac{x_k-y_k}{n(1+y_k)})$ for any strictly positive values of $x_1, ..., x_k, y_1, ..., y_k$ and hence realizing all (rational) multislopes in some open neighborhood of $\bar{0} \in \mathbb{R}^k$.

Lemma 3.8. *Suppose the weights $\bar{x}, \bar{y}$ are distinct and have strictly positive coordinates. Then each lamination $\lambda_{(\bar{x},\bar{y})}$, contains only noncompact leaves. Furthermore, each lamination $\lambda_{(\bar{x},\bar{y})}$ extends to a taut foliation $\mathcal{F}_{(\bar{x},\bar{y})}$, which realizes the same multislope.*

Proof. Suppose that $\lambda_{(\bar{x},\bar{y})}$ contains a compact leaf L. Such a leaf determines a transversely invariant measure on B given by counting intersections with L.

Now focus on any i, j, where $0 \leq i, j < n$. By considering, for example, a simple closed curve in F_i parallel to the arc α_i^j, we see that there is an oriented simple closed curve in F_i which intersects the branching locus of B_σ exactly k times that has orientation consistent with the branched locus. Since this is true for all possible i, j, it follows that the only transversely invariant measure B can support is the one with all weights on the branches D_i^j necessarily 0. But this means that $\lambda_{(\bar{x},\bar{y})}$ realizes multislope $\bar{0}$ and hence that $\bar{x} = \bar{y}$.

The complementary regions to the lamination $\lambda_{(\bar{x},\bar{y})}$ are product regions. Filling these up with product fibrations, we get the required foliation $\mathcal{F}_{(\bar{x},\bar{y})}$, which also has no compact leaves and is therefore taut.

□

REFERENCES

1. Danny Calegari and Nathan M. Dunfield, *Laminations and groups of homeomorphisms of the circle*, Invent. Math. **152** (2003), 149–207.
2. B. Casler, *An imbedding theorem for connected 3-manifolds with boundary*, Proc. A.M.S. **16** (1965), 559–566.
3. Yakov Eliashberg and William P. Thurston, *Confoliations*, University Lecture Series **13**, American Mathematical Society (1998).
4. David Gabai, *Foliations and the topology of 3-manifolds*, J. Differential Geom. **18** (1983), no. 3, 445–503.
5. David Gabai, *Foliations and the topology of 3-manifolds. II*, J. Differential Geom. **26** (1987), no. 3, 461–478.
6. David Gabai, *Foliations and the topology of 3-manifolds. III*, J. Differential Geom. **26** (1987), no. 3, 479–536.
7. D. Gabai and U. Oertel, *Essential Laminations in 3-Manifolds*, Ann. Math. **130**, (1989), 41–73.
8. Sylvain Gervais , *A finite presentation of the mapping class group of a punctured surface.*, Topology **40** (2001), no. 4, 703-725.
9. A. Haefliger, *Varietes feuillets*, Ann. Scuola Norm. Sup. Pisa (3) **16** (1962) 367–397
10. K. Honda, W.H. Kazez and G. Matic, *Right-veering diffeomorphisms of compact surfaces with boundary II*, Geometry & Topology, **12**, (2008), 2057–2094.
11. W. H. Kazez and R. Roberts, *Fractional Dehn twists in knot theory and contact topology*, `ArXiv:1201.5290`.
12. Tao Li, *Laminar branched surfaces in 3-manifolds*, Geom. Top. **6** (2002), 153–194.
13. Tao Li, *Boundary train tracks of laminar branched surfaces.*, Topology and geometry of manifolds (Athens, GA, 2001), 269-285, Proc. Sympos. Pure Math., **71**, Amer. Math. Soc., Providence, RI, 2003.
14. W.B. Lickorish, *A foliation for 3-Manifolds*, Ann. of Math. (2) **82** 414–420 1965.
15. S.P. Novikov, *The topology of foliations.* (Russian), Trudy Moskov. Mat. Ob. 14 1965 248–278.
16. U. Oertel, *Incompressible branched surfaces*, Invent. Math. **76** (1984), 385–410.
17. U. Oertel, *Measured laminations in 3-manifolds*, Trans, A.M.S. **305** (1988), 531–573.
18. Peter Ozsváth and Zoltán Szabó, *Holomorphic disks and genus bounds*, Geom. Topol. **8** (2004), 311–334.
19. Palmeira, Carlos Frederico Borges, *Open manifolds foliated by planes*, Ann. Math. (2) **107** (1978), no. 1, 109–131.
20. Rachel Roberts, *Taut foliations in punctured surface bundles. I*, Proc. London Math. Soc. (3) **82** (2001), no. 3, 747-768.
21. Rachel Roberts, *Taut foliations in punctured surface bundles, II*. Proc. London Math. Soc. (3) **83** (2001), no. 2, 443–471.
22. Rachel Roberts, John Shareshian and Melanie Stein, *Infinitely many hyperbolic 3-manifolds which contain no Reebless foliation*, J.A.M.S. **16(3)**, 639–379.

23. Georges Reeb, *Sur certaines propriétés topologiques des variétés feuilletées*, Actualités Sci. Indust. **1183** (1952), 91–158.
24. Dale Rolfsen, *Knot and Links*, Mathematics Lecture Series, No. 7. Publish or Perish, Inc., Berkeley, Calif., 1976.
25. H. Rosenberg, *Foliations by planes*, Topology **6** (1967) 131–138
26. William Thurston, *A norm for the homology of 3-manifolds*, Mem. Amer. Math. Soc. 59 (1986), no. 339, i-vi and 99-130.
27. William Thurston, *Three-manifolds, foliations and circles, II*, Unfinished manuscript, 1998.
28. R. Williams, *Expanding attractors*, Inst. Hautes Études Sci. Publ. Math. **43** (1974), 169–203.
29. John Wood, *Foliations on 3-manifolds*, Ann. of Math. **89(2)** (1969), 336–358.

MATHEMATICS DEPARTMENT, WASHINGTON UNIVERSITY IN ST LOUIS, ST LOUIS, MO 63130, USA